

TWO-DAY COLONOSCOPY INSTRUCTIONS: MIRALAX & SUPREP

Procedure Date: Provider Name	::
-------------------------------	----

The facility will call you with the instructions and arrival time 2 business days before your procedure.

FACILITY	ADDRESS	PHONE #
WELLNESS AVENUE SURGERY CENTER	2863 Wellness Ave. Orange City, FL	386-297-7239
CURRENCY CIRCLE SURGERY CENTER	835 Currency Circle, Suite 1021 Lake Mary, FL	407-544-6643
FHC PLANS SURGERY CENTER	2777 Enterprise Rd. Orange City, FL	386-481-6289
ADVENTHEALTH FISH MEMORIAL	1055 Saxon Blvd. Orange City, FL	386-917-5000
LAKE MONROE HOSPITAL	1401 West Seminole Blvd. Sanford FL	888-894-2106

Follow-up Appointment Date and Time:

 OFFICE	ADDRESS	PHONE #
ORANGE CITY	2884 Wellness Avenue, Suite 100 Orange City, FL	386-668-2221
LAKE MARY	835 Currency Circle, Suite 1001 Lake Mary, FL	407-749-6656
DELAND	929 N Spring Garden Ave, Suite 150 DeLand, FL	386-339-2692

INSTRUCTIONS

- Continue taking routine medications, including blood pressure and heart medications.
- If **diabetic**, take ½ of your usual dose of insulin the morning of your procedure.
- **Before stopping any blood thinner**, contact your prescribing provider.
- 7 days before your procedure, hold/do not take Mounjaro, Ozempic, Rybelsus, Trulicity, or Wegovy
- **5 days before** your procedure, **hold/do not take** the following medications:
 - Coumadin, Plavix, Effient, or Brilinta
 - Aleve, Motrin, Naproxen, or Ibuprofen [Tylenol is safe to use]
 - Iron supplements, vitamins, or herbal supplements
- 2 days before your procedure, hold/do not take Pradaxa, Eliquis, or Xarelto.
- **5 days before** the procedure, follow a low fiber diet.
 - Avoid: Raw vegetables, whole grains, nuts, seeds, corn, beans, oatmeal, and brown rice
 - Allowed foods (not limited to): Cooked or canned fruits & vegetables, white bread/rice/pasta, tender meats, and potatoes without skin
- You must have a friend or family member drive you to and from the facility. No form of taxi or ride-sharing service is permitted.

PURCHASE AT THE PHARMACY/GROCERY STORE

- Gatorade, 64-ounce bottle or Crystal Light (Recommended for diabetics) (No dark colors)
- Miralax, 238g bottle
- Dulcolax, 6 tablets
- Magnesium citrate, 10-ounce bottle
- Baby wipes or Desitin (Optional)

ACCEPTABLE CLEAR LIQUIDS

- **Soups:** Broth (chicken, beef, or vegetable)
- **Juices:** White grape juice, apple juice, etc.
- **Desserts:** Jell-O, Italian ice, Popsicles, etc.
- Sports Drinks: Powerade, Gatorade, Propel
- Beverages: Tea, Kool-aide, water, black coffee



TWO DAYS BEFORE PROCEDURE - DULCOLAX AND MIRALAX

- Do Not Eat Any Solid Food the Entire Day Before the Procedure.
- Have a clear liquid diet throughout the day. Avoid liquids that are red or purple in color such as red grape or cranberry juice. Avoid ALL dairy products and juices with pulp such as orange or grapefruit juice.
- 1. At **5:00 PM**, take 4 Dulcolax tablets with water.
- 2. At **5:30 PM**, drink the entire 10-ounce bottle of Magnesium citrate.
- 3. At **6:00 PM,** mix the entire 238g bottle of Miralax with the 64 oz bottle of Gatorade or Crystal Light in a pitcher. Stir the solution until the Miralax powder completely dissolves. Drink an 8-ounce glass every 10-15 minutes until the solution is gone (Drinking this with a straw helps).
- 4. At 8:00 PM, take 2 Dulcolax tablets with water.
- 5. At **9:00 PM**, continue to drink an 8-ounce solution of the Miralax every 10-15 minutes until the rest of the solution is consumed.

DAY BEFORE PROCEDURE - SUPREP

- Do Not Eat Any Solid Food the Entire Day Before the Procedure.
- Have a clear liquid diet throughout the day. Avoid liquids that are red or purple in color such as red grape or cranberry juice. Avoid ALL dairy products and juices with pulp such as orange or grapefruit juice.
- Until **3 hours before** your procedure time, you may drink <u>clear liquids</u> (NO RED/DARK COLORS). You may also take any necessary medication with a small sip of water. At the **3-hour mark**, take nothing by mouth including water.

Ify	our procedure is before 12:00 PM:	If your procedure is After 12:00 PM:	
1.	At 5:00 PM , pour (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add cold water to the 16-ounce line and drink all the liquid in the cup.	1. At 7:00 PM , pour (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add cold water to the 16-ounce line and drink all the liquid in the cup.	
2.	Drink (2) more 16-ounce cups of water (or clear liquid) over the next 1 hour.	2. Drink (2) more 16-ounce cups of water (or clear liquid) over the next 1 hour.	
3.	At 11:00 PM , pour (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add cold water to the 16-ounce line and drink all the liquid in the cup.	3. At 6:00 AM , pour (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add cold water to the 16-ounce line and drink all the liquid in the cup.	
4.	Drink (2) more 16-ounce cups of water (or clear liquid) over the next 1 hour.	4. Drink (2) more 16-ounce cups of water (or clear liquid) over the next 1 hour.	

DAY OF PROCEDURE

- Do Not Eat Any Solid Food the Day of the Procedure.
- Until **3 hours before** your procedure time, you may drink <u>clear liquids</u> (NO RED/DARK COLORS). You may also take any necessary **medication** with a small sip of water. At the **3-hour mark**, take nothing by mouth including water.



CANCELLATION FEES

#